

NATtrol™ Parainfluenza Virus Control

PRODUCT DESCRIPTION:

NATtrol™ Parainfluenza Virus Control contains purified, intact viral particles that have been chemically modified to render them noninfectious and refrigerator stable. The control is supplied in a proprietary matrix.

INTENDED USE:

NATtrol™ Parainfluenza Virus Control is designed to evaluate the performance of nucleic acid tests for determination of the presence of Parainfluenza Types 1, 2, 3 and 4B nucleic acid. NATPIV-ERC enables laboratories to monitor test variation, lot-to-lot test kit performance, operator variation, and can provide assistance in identifying random or systemic error.

MATERIALS SUPPLIED:

Each control pack contains 6 x 0.5 mL vials.

WARNINGS AND PRECAUTIONS:

- NATtrol™ inactivation was carried out on the viral stocks used to formulate the control. The inactivation was verified in a standard microbiological growth protocol.
- This product contains inactivated viruses and materials of human and animal origin. Safe practices suggest that the controls be considered potentially infectious and to use Universal Precautions when handling.
- Refer to CDC guidelines and local regulations for handling and disposal.
- The matrix used in the manufacture of this product is treated with 0.09% sodium azide. It was manufactured from Human Serum Albumin that has been tested and found to be non-reactive at the donor level for HIV-1/HIV-2 Antibody, HBsAg and HCV Antibody by FDA licensed donor screening test methods. All materials are also tested for HIV-1 and HCV by FDA approved Nucleic Acid Test (NAT) methods.
- Heat inactivated Fetal Bovine Serum used in the manufacture of this product meets applicable USDA requirements for abattoir sourced animals, traceability, and country of origin. The materials were collected at USDA licensed establishments or legally imported from countries recognized by the USDA as negligible or controlled for risk for Bovine Spongiform Encephalopathy (BSE) and other exotic disease agents. Donor animals were inspected ante and postmortem at the abattoir as required by the USDA.
- Do not use past the expiration date on the label.
- To avoid cross-contamination, use separate pipette tips for all materials.

RECOMMENDED STORAGE:

NATtrol™ Parainfluenza Virus Control should be stored at 2-8°C.

INSTRUCTIONS FOR USE:

- Mix tube vigorously for at least 5 secs.
- Process according to manufacturer's instructions for sample to result assays.
- Extract nucleic acid prior to use in downstream assays that are not sample to result.

LIMITATIONS:

- FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES.
- Quality control materials should be used in accordance with local, state, federal, and accreditation requirements.
- This product is not intended to replace the manufacturer's controls provided with the assay.

PINATPIV-ERC Revision: 03

PCA# 24-292 Effective Date: 09/18/2024 Page 1 of 2



NATtrol™ Parainfluenza Virus Control

EXPECTED RESULTS:

- The expected results shown in Table 1 below are tested using ZeptoMetrix in-house real-time PCR assay.
- It is the responsibility of the end user to establish their own acceptance criteria for the control using their laboratory's procedures.

Table 1

Catalog Number	Virus	Expected Result
NATPIV-ERC	Parainfluenza Type 1	DETECTED1
	Parainfluenza Type 2	DETECTED ²
	Parainfluenza Type 3	DETECTED ²
	Parainfluenza Type 4B	DETECTED ³

¹Based on in-house real time PCR assay targeting HN gene region.

Symbols used in the labeling of this product:

ш	Manufacturer	ł.	Temperature Limitation
RUO	For Research Use Only	₽	Expiration Date
REF	Catalog Number	⊗	Biological Risk
LOT	Batch Code		

PINATPIV-ERC Revision: 03 Effective Date: 09/18/20

²Based on in-house real time PCR assay targeting nucleocapsid protein (NP) gene region.

³Based on in-house real time PCR assay targeting phosphoprotein (P) gene region.